

Treating the enlarged prostate with prostate artery embolization (PAE)

- PAE treats urinary symptoms while preserving much of the prostate
- PAE is minimally invasive with a quick recovery period
- With PAE, greater than 84% of patients see significant improvement

Prostate artery embolization (PAE) is an emerging treatment for men with benign prostatic hyperplasia (BPH), also known as an enlarged prostate.

Half of men age 50 and older and 80% of men age 80 or older show signs of enlarged prostate.

Overgrowth of the prostate tissue can obstruct the urethra, leading to discomfort and difficulty urinating. PAE is especially promising for patients that have failed medication therapy or surgery, or prefer to avoid surgery.

Take a survey to find out if you have an enlarged prostate at ausrad.com.

ARA CONVENIENCE

- Exceptional patient care
- Most insurance plans accepted and filed
- Flexible office hours
- Handicapped-accessible parking

AUSTIN CENTER BOULEVARD & ACB BREAST IMAGING
6818 Austin Center Blvd., Suite 101
Austin, TX 78731
(512) 795-8505

CEDAR PARK & CEDAR PARK BREAST IMAGING
12800 W. Parmer Ln., Suite 200
Cedar Park, TX 78613
(512) 485-7199

DRIPPING SPRINGS
170 Benney Lane, Suite 101
Dripping Springs, TX 78620
(512) 776-1176

GEORGETOWN
3201 S. Austin Avenue, Suite 105
Georgetown, TX 78626
(512) 863-4648 or
(512) 519-3441

KYLE & KYLE BREAST IMAGING CENTER
4211 Benner Rd., Suite 100
Kyle, TX 78640
(512) 776-1150

MANOR
12700 Lexington St., Suite 300
Manor, TX 78653
(512) 776-1158

MEDICAL PARK TOWER
1301 W. 38th Street, Suite 118
Austin, TX 78705
(512) 454-7380

★ **MIDTOWN**
901 W. 38th Street, Suite 100
Austin, TX 78705
(512) 519-3456

MUELLER & MUELLER CHILDREN'S IMAGING
1301 Barbara Jordan Blvd.
Suite 104
Austin, TX 78723
(512) 480-0761

QUARRY LAKE
4515 Seton Center Parkway
Suite 105
Austin, TX 78759
(512) 519-3402

ROCK CREEK PLAZA
2120 N. Mays, Suite 220
Round Rock, TX 78664
(512) 238-7195

SAN MARCOS
1348 B Texas 123 South
San Marcos, TX 78666
(512) 392-1831 or
(888) 261-2149

SOUTHWEST MEDICAL VILLAGE
5625 Eiger Road, Suite 165
Austin, TX 78735
(512) 519-3475

SOUTHWOOD
1701 W. Ben White Blvd., Suite 170
Austin, TX 78704
(512) 428-9090

WESTLAKE
5656 Bee Caves Road
Building H, Suite 200
Austin, TX 78746
(512) 328-4984

WILLIAM CANNON
2501 W. William Cannon Drive
Building 5
Austin, TX 78745
(512) 346-7311

WILSON PARKE
11714 Wilson Parke Ave., Suite 175
Austin, TX 78726
(512) 519-3457

★ Locations offering PAE

Are you challenged with frequent urination?

You may have an enlarged prostate. Prostate artery embolization (PAE) can help while preserving the prostate.




OCT 2023

SCHEDULING
(512) 467-9729

FAX REFERRALS
(512) 343-9099





What are the symptoms of enlarged prostate?

- Frequent urination at night
- Urination that produces a small amount of urine
- Hesitant or interrupted urine stream
- Leaking or dribbling urine
- Sudden and urgent need to urinate
- Feeling like the bladder is not completely emptied after urinating
- Occasional pain when urinating
- Not being able to urinate at all due to obstruction

How is prostate artery embolization (PAE) done?

In PAE, a catheter is inserted into the femoral artery in the groin and guided through imaging to the prostate artery on both sides of the enlarged gland. Once positioned next to the prostate, microscopic beads are released into the artery to block blood flow, causing the prostate to shrink over time. The benefit to the patient is immense because PAE avoids the surgical practice of removing large parts of the prostate, which can result in a longer recovery period and sexual and urological dysfunction.

What steps are taken to have the PAE procedure?

You will need to visit a urologist and obtain a referral to the ARA interventional clinic. At ARA, you will fill out a short questionnaire, provide your health history

and get a physical examination to evaluate candidacy for PAE. If it's determined that PAE is a good fit, you will meet with an ARA interventional radiologist who will discuss treatment options and if pre-procedure tests are needed. You may need a CT study of the pelvis or other imaging.

What is the procedure like?

The procedure is done on an outpatient basis at ARA's Midtown location. PAE takes about 1.5 hours and patient can generally leave the clinic about 3 hours afterward. There is no need for general anesthesia. PAE is not considered painful and some patients elect to have no sedation. If you wish, you will be given a sedative medication like that given for a colonoscopy, which has the effect referred to as "twilight sleep."

What is recovery like?

There may be mild pain or discomfort after the procedure and a week of home recovery time is recommended.

What are the results of PAE?


Greater than 84 percent of patients experience significant improvement. Rarely, symptoms return and, if so, PAE can be repeated.

Does insurance cover PAE?


After your consultation, if you require insurance pre-certification for the procedure from your insurance provider, ARA will assist you. Generally, insurance companies will pay for prostate embolization but each plan is different, so check with your provider.

Take a survey about your symptoms at ausrad.com/pae. Talk to your urologist about PAE and if it might be the right procedure for you. You can schedule an ARA consultation at (512) 453-6100.


How Prostate Artery Embolization Works



1. When the prostate is enlarged, as shown, it constricts the urethra making urination difficult. PAE is a minimally-invasive way of reducing the size of the prostate.



2. The procedure begins with the insertion of a catheter into the an artery at the groin or wrist. The catheter is advanced through the blood vessels until reaching the arteries of the prostate.



3 & 4. Small beads, typically made of nonreactive gelatin, are released through the catheter into the small blood vessels that supply prostate, blocking the flow of blood. The beads are between 100 and 500 micrometers in diameter. They do not disintegrate, but remain permanently in place and cannot be felt by the patient. This procedure is done on both sides of the prostate. Remaining vessels supply sufficient blood to maintain a healthy prostate.



5. With a lowered blood supply, the prostate shrinks over time and there is less pressure on the urethra, allowing urine to flow freely.