



DIAGNOSTIC IMAGING MRI PROSTATE ORDER FORM

Patient name: _____ Phone: _____

Diagnosis: _____

PSA (prostate-specific antigen) value: _____

- ☐ MRI Prostate with perfusion with and without contrast (72197— MRI Pelvis with and without contrast)
- ☐ MRI Prostate post prostatectomy with and without contrast (72197—MRI Pelvis with and without contrast)
- ☐ MRI Prostate radiation treatment planning (72195— MRI Pelvis without contrast)

Provider scheduling line: 512-458-9098 | Scheduling fax: 512-836-8869

Appointment date: _____ Appointment time: _____ Arrival time: _____

Ordering physician: _____ Signature: _____ Date: _____

Appointment location:

- ☐ **ARA Cedar Park** – 12800 W. Parmer Ln, Suite 200, Cedar Park, TX 78613
- ☐ **ARA Georgetown** – 3201 S. Austin Ave., Suite 105, Georgetown, TX 78626
- ☐ **ARA Kyle** – 4211 Benner Rd., Suite 100, Kyle, TX 78640
- ☐ **ARA Medical Park Tower** – 1301 W. 38th St., Suite 118, Austin, TX 78705
- ☐ **ARA San Marcos** – 1348 Texas 123 South, Ste B, San Marcos, TX 78666
- ☐ **ARA Southwood** – 1701 W. Ben White Blvd., Suite 170, Austin, TX 78704
- ☐ **ARA Westlake** – 5656 Bee Caves Road, Bldg. H, Suite 200, Austin, TX 78746
- ☐ **ARA William Cannon** – 2501 W William Cannon Drive, Bldg. 5, Austin, TX 78745

Prep instructions for MRI Prostate with perfusion and MRI Prostate post prostatectomy:

Does not apply to prostate radiation treatment planning (72195).

General instructions

- Arrive 30 minutes prior to exam.
- Please leave all jewelry and metal objects at home.
- If applicable, bring an identification card for any implant devices in the body, or any handheld or remote programmer device.
- You must wait 2 weeks after any iron infusion before having this exam.
- Morning appointments preferred.
- Children 10 and under must be supervised at all times and may not accompany patients beyond the lobby.

Prep instructions

- **Abstain from ejaculation for 3 days** prior to this exam.
- **12 to 16 hours prior** to the exam, take 2 Dulcolax tablets.
- **8 hours before the exam**, have your last meal. Limit to a small sandwich, Jell-O, soup, or broth. No dairy products. Do not eat anything else before the exam. You may continue to drink clear liquids (water).
- **1 hour prior** to leaving home/arriving at the clinic, perform a water enema. Additional information is available at <https://www.ausrad.com/wp-content/uploads/2025/04/ARA-Instructions-for-Tap-Water-Enema.pdf>
- **Patients on dialysis** need to coordinate their next dialysis appointment for within 24 hours after this exam to help move the contrast out of the body. Contact your physician for additional information.